

WIN 2 / 2014 IE

**PATENTS** 

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

)
)
) Art Ünit: <b>1614</b>
)
) Examiner: Unassigned
)
)
)

## INFORMATION DISCLOSURE STATEMENT

Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450

Sir:

The citation of information on the attached Form PTO/SB/08 is made pursuant to 37 C.F.R. §§ 1.56, 1.97, and 1.98. A copy of each cited item is enclosed.

The citation of this information does not constitute an admission of priority or that any cited item is available as a reference, or a waiver of any right the applicant may have under applicable statutes, Rules of Practice in patent cases, or otherwise.

Respectfully submitted,

hn K. McDonald, Ph.D.

Reg. No. 42,860

KILPATRICK STOCKTON LLP

1100 Peachtree Street, Suite 2800 Atlanta, Georgia 30309-4530

(404) 815-6500

Our Docket: 05501-0202 (43150/287577)

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, on May 19, 2004.

John K. McDonald, Ph.D. - Reg. No. 42,860

PTO/SB/08A (08-00)

Approved for use through 10/31/2002 OMB 0651-0031

U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE Under the Paperwork Authors, no persons are required to respond to a collection of information unless it contains a valid OMB control number

Substitute for Form 1449/A/PTO

1

Sheet

## INFORMATION DISCLOSURE STATEMENT BY APPLICANT

(use as many sheets as necessary

of

WO

4.

2

02/081703

Cor	nplete if Known
Application Number	10/621,113
Filing Date	July 16, 2003
First Named Inventor	J. David Lambeth et al.
Group Art Unit	1614
Examiner Name	Unknown
Attorney Docket Number	43150/287577 (05501-0202)

10-17-2002

**U.S. PATENT DOCUMENTS** U.S. Patent Document Date of Publication of Pages, Columns, Lines, Where Kind Code<sup>2</sup> Name of Patentee or Applicant Cited Document Relevant Passages or Relevant Number Cite No.1 Examiner of Cited Document MM-DD-YYYY Figures Appear (if known) Initials 6,620,603 LAMBETH et al. 09-16-2003 1. **FOREIGN PATENT DOCUMENTS** Date of Publication Foreign Patent Document Pages, Columns, Lines, Examiner Name of Patentee or Kind Code<sup>5</sup> of Where Cite Initials Cited Document Office<sup>3</sup> Number<sup>4</sup> Applicant of Cited Document Relevant Passages or (if known) MM-DD-YYYY Relevant Figures Appear WO 05/18/2000 2. 00/28031 A2 **Emory University** WO 01/087957 11-22-2001 3. **A3 Emory University** 

**OTHER INFORMATION - NON PATENT LITERATURE DOCUMENTS** 

**Emory University** 

Examiner Initials	Cite No.1	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published	T <sup>2</sup>			
	5.	Burdon, R.H., "Superoxide and hydrogen peroxide in relation to mammalian cell				
	_	proliferation," Free Radical Biol. Med., Volume 18, No. 4, pp. 775-794 (1995).	<u> </u>			
•	6.	Cheng, G., et al., Gene, 269: pp. 131-140, May 16, 2001.	<u> </u>			
	7.	Church, S.L., et al., "Increased manganese superoxide dismutase expression suppresses the				
		malignant phenotype of human melanoma cells," Proc. Natl. Acad. Sci. USA, Volume 90,	İ			
		pp. 3113-3117 (1993).				
	8.	······································				
		multidomain oxidase/peroxidase with homology to the phagocyte oxidase subunit				
		gp91phox," J. Cell Biol., 154(4): pp. 879-91, August 20, 2001.				
	9.					
		an increase in superoxide dismutase activity in a revertant subclone of sarcoma virus-				
		infected mammalian cells," Can. Res., Volume 42, pp. 609-617 (1982).				
	10.	<sup>0.</sup> Fukui, T., et al., "p22phox mRNA expression and NADPH oxidase activity are increased in				
		aortas from hypertensive rats," Circ. Res., Volume 80, No. 1, pp. 45-51 (1997).				
	11.	Griendling, K. K., et al., "Angiotensin II stimulates NADH and NADPH oxidase activity in				
		cultured vascular smooth muscle cells," Circ. Res., Vol. 74, No. 6, pp. 1141-1148 (1994).				
	12.	Irani, K., et al., "Mitogenic signaling mediated by oxidants in ras-transformed fibroblasts,"	"			
		Science, Vol. 275, No. 5306, pp. 1649-1652 (1997).				
	13.	Lambeth, J. D., et al., "Novel homologs of gp91phox," Trends Biochem. Sci., (10): pp. 459-	9-			
		61, October 25, 2000.				
	14.					
		for detecting superoxide anion radical production by enzymatic and cellular systems," J.				
		Biol. Chem., Vol. 273, No. 4, pp. 2015-2023 (1998).				
	15.	Matsubara, T., et al., "Increased superoxide anion release from human endothelial cells in				
		response to cytokines," J. Immun., Vol. 137, No. 10, pp. 3295-3298 (1986).				
1	<del></del>	35 1 000 11 11 11 11 11 11 11 11 11 11 11	<del>.                                      </del>			

<sup>1</sup>Unique citation designation number. <sup>2</sup>See attached Kinds of U.S. Patent Documents. <sup>3</sup>Enter Office that issued the document, by the two-letter code (WIPO Standard ST.3). <sup>4</sup>For Japanese patent document, the indication of the year of the reign of the Emperor must precede the serial number of the patent document. <sup>5</sup>Kind of document by the appropriate symbols as indicated on the document under WIPO Standard ST. 16 if possible. <sup>6</sup>Applicant is to place a check mark here if English language translation is attached.

Substitute for Form 1448/APTO   Substitute for Form 1448/APTO   STATEMENT BY APPLICANT	•	<b>`</b>		U.S. Patent and Trade	PTO/SB/08B (0 proved for use through 10/31/2002 OMB 0651 mark Office: U.S. DEPARTMENT OF COMMI	-0031		
Application Number   10/621,113   Filing Date   July 16, 2003	Under the i	nder the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number  Complete if Known						
Filing Date   July 16, 2003   First Named Inventor   J. David Lambeth et al.   Group Art Unit   Learning   Unknown   Attorney Docket Number   43150/287577 (05501-0202)	Substitute for Form 1449/A/PTO							
STATEMENT BY APPLICANT   Group Art Unit   Info	INFORMATION DISCLOSURE		RMATION DISCLOSURE	Filing Date				
Group Art Unit   Examiner Name   Unknown				First Named Inventor				
Sheet   2   of   2   Attorney Docket Number   43150/287577 (05501-0202)		JIA		Group Art Unit	1614			
Cite   Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, calatog, etc.), data, page(s), volume-issue number(s).   17				Examiner Name	Unknown			
Cite   Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), data, page(s), volume-issue number(s). publisher, city and/or country where published  16.   Meier, B., et al., "Human fibroblasts release reactive oxygen species in response to interleukin-1 or tumor necrosis factor-α," Biochem. J., Vol. 263, No. 2, pp. 539-545 (1989).  17.   Pagano, P. J., et al., "Localization of a constitutively active, phagocyte-like NADPH oxidase in rabbit aortic adventitia: Enhancement by angiotensin II," Proc. Natl. Acad. Sci. USA, Volume 94, No. 26, pp. 14483-14488 (1997).  18.   Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-κB," Chem. & Biol., Vol. 2, No. 1, pp. 13-22 (1995).  19.   Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-κB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).  20.   Sundaresan, M., et al., "Requirement for generation of H <sub>2</sub> O <sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).  21.   Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22.   Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23.   Ushio-Fukai M., et al., "p22 <sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24.   Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-28	Sheet	2	of 2	Attorney Docket Number	43150/287577 (05501-0202)	)		
Cite   Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), data, page(s), volume-issue number(s). publisher, city and/or country where published  16.   Meier, B., et al., "Human fibroblasts release reactive oxygen species in response to interleukin-1 or tumor necrosis factor-α," Biochem. J., Vol. 263, No. 2, pp. 539-545 (1989).  17.   Pagano, P. J., et al., "Localization of a constitutively active, phagocyte-like NADPH oxidase in rabbit aortic adventitia: Enhancement by angiotensin II," Proc. Natl. Acad. Sci. USA, Volume 94, No. 26, pp. 14483-14488 (1997).  18.   Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-κB," Chem. & Biol., Vol. 2, No. 1, pp. 13-22 (1995).  19.   Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-κB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).  20.   Sundaresan, M., et al., "Requirement for generation of H <sub>2</sub> O <sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).  21.   Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22.   Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23.   Ushio-Fukai M., et al., "p22 <sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24.   Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-28			OTHER INCORNATION, NON B	ATENT LITERATURE DOCUME	INTE			
16. Meier, B., et al., "Human fibroblasts release reactive oxygen species in response to interleukin-1 or tumor necrosis factor-\(\alpha\)," Biochem. J., Vol. 263, No. 2, pp. 539-545 (1989).  17. Pagano, P. J., et al., "Localization of a constitutively active, phagocyte-like NADPH oxidase in rabbit aortic adventitia: Enhancement by angiotensin II," Proc. Natl. Acad. Sci. USA, Volume 94, No. 26, pp. 14483-14488 (1997).  18. Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-kB," Chem. & Biol., Vol. 2, No. 1, pp. 13-22 (1995).  19. Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).  20. Sundaresan, M., et al., "Requirement for generation of H2O2 for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).  21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23. Ushio-Fukai M., et al., "p22 <sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).			Include name of the author (in CAPITAL	LETTERS), title of the article (when a	ppropriate), title of the			
<ol> <li>Meier, B., et al., "Human fibroblasts release reactive oxygen species in response to interleukin-1 or tumor necrosis factor-α," Biochem. J., Vol. 263, No. 2, pp. 539-545 (1989).</li> <li>Pagano, P. J., et al., "Localization of a constitutively active, phagocyte-like NADPH oxidase in rabbit aortic adventitia: Enhancement by angiotensin II," Proc. Natl. Acad. Sci. USA, Volume 94, No. 26, pp. 14483-14488 (1997).</li> <li>Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-κB," Chem. &amp; Biol., Vol. 2, No. 1, pp. 13-22 (1995).</li> <li>Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-κB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).</li> <li>Sundaresan, M., et al., "Requirement for generation of H<sub>2</sub>O<sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).</li> <li>Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ol>					volume-issue number(s),	T'		
<ol> <li>Pagano, P. J., et al., "Localization of a constitutively active, phagocyte-like NADPH oxidase in rabbit aortic adventitia: Enhancement by angiotensin II," Proc. Natl. Acad. Sci. USA, Volume 94, No. 26, pp. 14483-14488 (1997).</li> <li>Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-κB," Chem. &amp; Biol., Vol. 2, No. 1, pp. 13-22 (1995).</li> <li>Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-κB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).</li> <li>Sundaresan, M., et al., "Requirement for generation of H<sub>2</sub>O<sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).</li> <li>Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>Ushio-Fukai M., et al., "p22phox is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ol>		16.			gen species in response to			
in rabbit aortic adventitia: Enhancement by angiotensin II," Proc. Natl. Acad. Sci. USA, Volume 94, No. 26, pp. 14483-14488 (1997).  18. Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-kB," Chem. & Biol., Vol. 2, No. 1, pp. 13-22 (1995).  19. Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).  20. Sundaresan, M., et al., "Requirement for generation of H <sub>2</sub> O <sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).  21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23. Ushio-Fukai M., et al., "p22phox is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.			interleukin-1 or tumor necrosis factor-α,	"Biochem. J., Vol. 263,	No. 2, pp. 539-545 (1989).			
in rabbit aortic adventitia: Enhancement by angiotensin II," Proc. Natl. Acad. Sci. USA, Volume 94, No. 26, pp. 14483-14488 (1997).  18. Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-kB," Chem. & Biol., Vol. 2, No. 1, pp. 13-22 (1995).  19. Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).  20. Sundaresan, M., et al., "Requirement for generation of H <sub>2</sub> O <sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).  21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23. Ushio-Fukai M., et al., "p22phox is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.		17.	Pagano, P. J., et al., "Localization of a c	onstitutively active, pha	gocyte-like NADPH oxidase			
<ol> <li>Schmidt, K. N., et al., "The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-κB," Chem. &amp; Biol., Vol. 2, No. 1, pp. 13-22 (1995).</li> <li>Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-κB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).</li> <li>Sundaresan, M., et al., "Requirement for generation of H<sub>2</sub>O<sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).</li> <li>Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ol>								
<ul> <li>activation of transcription factor NF-κB," Chem. &amp; Biol., Vol. 2, No. 1, pp. 13-22 (1995).</li> <li>19. Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-κB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).</li> <li>20. Sundaresan, M., et al., "Requirement for generation of H<sub>2</sub>O<sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).</li> <li>21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>23. Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b558," J. Biol.</li> </ul>								
<ol> <li>Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-κB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).</li> <li>Sundaresan, M., et al., "Requirement for generation of H<sub>2</sub>O<sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).</li> <li>Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b558," J. Biol.</li> </ol>		18.	Schmidt, K. N., et al., "The roles of hyd	lrogen peroxide and sup-	eroxide as messengers in the			
the activation of the NF-kB transcription factor and HIV-1," EMBO J., Vol. 10, No. 8, pp. 2247-2258 (1991).  20. Sundaresan, M., et al., "Requirement for generation of H <sub>2</sub> O <sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).  21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23. Ushio-Fukai M., et al., "p22 <sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.			activation of transcription factor NF-кВ,	"Chem. & Biol., Vol. 2,	, No. 1, pp. 13-22 (1995).			
<ul> <li>2247-2258 (1991).</li> <li>20. Sundaresan, M., et al., "Requirement for generation of H<sub>2</sub>O<sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).</li> <li>21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>23. Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ul>		19.	Schreck, R., et al., "Reactive oxygen intermediates as apparently widely used messengers in					
<ul> <li>20. Sundaresan, M., et al., "Requirement for generation of H<sub>2</sub>O<sub>2</sub> for platelet-derived growth factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).</li> <li>21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>23. Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ul>				n factor and HIV-1," El	MBO J., Vol. 10, No. 8, pp.			
factor signal transduction," Science, Vol. 270, pp. 296-299 (1995).  21. Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23. Ushio-Fukai M., et al., "p22 <sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b558," J. Biol.								
<ul> <li>Szatrowski, T.P., et al., "Production of large amounts of hydrogen peroxide by human tumor cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).</li> <li>Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ul>		20.		<b>—</b>	-			
cells," Canc. Res., Vol. 51, No. 3, pp. 794-798 (1991).  22. Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23. Ushio-Fukai M., et al., "p22 <sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b558," J. Biol.		21						
<ul> <li>Uhlinger, D.J., "Nucleoside triphosphate requirements for superoxide generation and phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).</li> <li>Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b558," J. Biol.</li> </ul>								
phosphorylation in a cell-free system from human neutrophils," Vol. 266, No. 31, pp. 20990-20997 (1991).  23. Ushio-Fukai M., et al., "p22 <sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.		22						
<ul> <li>23. Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ul>		22.						
<ul> <li>Ushio-Fukai M., et al., "p22<sup>phox</sup> is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).</li> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ul>				om numan neutrophiis,	voi. 200, No. 31, pp. 20990-			
NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.		23		critical component of	f the superavide generating			
vascular smooth muscle cells," J. Biol. Chem., Vol. 271, No. 38, pp. 23317-23321 (1996).  24. Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.		25.						
<ul> <li>Yan, T., et al., "Manganese-containing superoxide dismutase overexpression causes phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).</li> <li>Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b<sub>558</sub>," J. Biol.</li> </ul>			l	-	V			
phenotypic reversion in SV40-transformed human lung fibroblasts," Canc. Res., Vol. 56, pp. 2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.		24.						
2864-2871 (1996).  25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.			_					
25. Yu, L., et al., Biosynthesis of the phagocyte NADPH oxidase cytochrome b <sub>558</sub> ," J. Biol.					, , , , , , , , , , , , , , , , , , ,			
		25.		gocyte NADPH oxidase	e cytochrome b <sub>558</sub> ," J. Biol.			
			Chem., Vol. 272, No. 43, pp. 27288-272	94 (1997).				

Examiner Date Signature Considered
------------------------------------

<sup>&</sup>lt;sup>1</sup>Unique citation designation number. <sup>2</sup>Applicant is to place a check mark here if English language translation is attached.